

DIRECTIONS FOR USE
BEFORE TREATMENT, THE GRAIN SHOULD BE DRY
PREFERABLY WITH A MOISTURE CONTENT OF
LESS THAN 14% for effective pest control with
ACTELLC SUPER FOR BEST RESULTS GRAIN
SHOULD BE TREATED IMMEDIATELY AFTER
SHELLING AT THE TIME OF STORING. Established
infestations are difficult to control.
Good coverage is important though the fumigant action
partially compensates for this, Treated grain is protected
for at least six months.

RATES FOR USE
Cob-maize
It is not recommended to use ACTELLC SUPER on cob
maize to control the Larger Grain Borer. Better control of
the pests is obtained on shelled maize.

Shelled grain (maize, wheat, beans)
Mix 50 g (fill the provided measure to the mark) for every
50kg bag of grain using a shovel. The grain and
ACTELLC SUPER have to be thoroughly admixed for
effective control of larger grain borer. Sprinkle some
ACTELLC SUPER on the outside of the bag to reduce
re-infestation.

Spillage
Clean the contaminated surface using soil, sawdust or a
binding agent and bury in a safe place away from water
sources, and human habitation.

HANDLING PRECAUTIONS
SAFETY PRECAUTIONS
ACTELLC SUPER contains pirimiphos-methyl, which is
an organophosphate. Do not use the product if a doctor
has advised you not to work with organophosphate,
carbamate or thiocarbamate compounds. Do not smoke,
drink or eat while using this product. Do not breathe dust.
Avoid contact with skin and eyes.

For Safety when mixing
Dust may irritate your eyes. If you get any in your eyes,
wash away at once with plenty of clean water. Wear
protective clothing, hand gloves and a mask covering
nose and mouth.

For Safety after mixing
Brush any dust from your skin. Wash with soap and water.
Change clothing and wash work clothes. Burn or bury
empty packs, away from water sources and human
habitation. Do not re-use them for any other purpose. To
avoid harming fish, do not dump unwanted ACTELLC
SUPER in water.

Storage
Store in original tightly sealed satchets, away from children,
food and livestock in a cool, well-ventilated, dry place.
Stored under these conditions this product has a shelf life of
18 months.

Disposal
Destroy empty satchets by perforating, burning or burying
away from water sources. Do not re-use empty packs for
any other purpose. Dilute any spilled dust with wet sawdust,
sweep and bury away from water sources and human
habitation.

Environmental hazard
Toxic to aquatic organisms, Do not throwaway any dust or
containers into water sources.

FIRST AID INSTRUCTIONS
Inhalation: ACTELLC SUPER can cause sneezing and
coughing if it enters the respiratory system. Affected
persons should be moved to fresh air immediately.
Skin contact: Contaminated skin should be washed with
plenty of water and soap.
Eye contact: Hold eyelids apart and pour in a gentle
stream of clean water for 15 minutes. See a doctor.
Ingestion: If ACTELLC SUPER is swallowed, do not
induce vomiting. Take patient and this label to a doctor
immediately. If gastric lavage is performed, take care to
prevent aspiration of gastric contents. Consider
administration of activated charcoal and a laxative. Treat
symptomatically.

TOXICOLOGICAL INFORMATION
Symptoms of poisoning
Nausea, headache, loss of joint co-ordination; blurred
vision, stomach cramp and tightness in the chest. Severe
exposure may cause difficulties in breathing; in this case
oxygen and artificial respiration may be required.

Antidote: Atropine, in addition oxime preparations, under
medical supervision.

Notes to the physician
If poisoning is suspected, Atropine should be administered
immediately at 2-5 mg as an intramuscular injection
followed by 2 mg every 30 minutes until atropinization
occurs. Atropine must not be given to cyanosed patients;
administer oxygen first. Do not use opiates or barbiturates.
If convulsions occur administer diazepam (10 g
intravenously). Cholinesterase reactivators (pralidoxime)
should be given at the same time as atropine. Mild
poisoning: 1 g in 2-3 ml water (intramuscularly);
severe poisoning: 2 g in 30 ml water (intravenously).
In case of an emergency, call Toll free Nos: 0800720021
and 0800730030 (24hrs)

NOTICE TO USER
This pest control product is to be used only in accordance
with the directions on this label. It is an offence under the
Pest Control Products Act to use or store a pest control
product under unsafe conditions.

WARRANTY
Seller's guarantee is limited to the terms set out on the
label and subject thereto, the buyer assumes the risk to
persons or property arising from the use or handling of this
product and accepts the product on that condition.

MEMBER OF
AGROCHEMICALS
ASSOCIATION OF KENYA
REGISTRATION NUMBER 102

ACTELLC SUPPER

Dusting Powder


**Insecticidal dust for the control of
stored maize and small grain cereals
pests including control of larger
grain borer,**

**READ THE LABEL BEFORE USING
COMMERCIAL AND AGRICULTURAL CLASS
KEEP LOCKED OUT OF REACH OF
CHILDREN**

**GUARANTEE : Pirimiphosmethyl
(DHAMANA) 16g/kg + Permethrin 3g/kg.
REGISTRATION No.: PCPB(CR) 0150**

**Dawa ya kuzuia dudumuzi
pamoja na wadudu waharibifu
wa nafaka.**

**SOMA MAELEZO KABLA YA KUTUMIA
DAWA HII.
SEHEMU YA BIAHARA NA KILIMO
HIFADHI MBALI NA WATOTO
NAMBARI YA USAJILI: PCPB(CR) 0150**



**MANUFACTURER: Syngenta Crop Protection AG,
Basle, Switzerland**

DATE OF MANUFACTURE (TAREHE YA KUTEN- GENEZWA)	BATCH NO.: (NA NAMBARI YA KIFURUSHI)

**SHELF LIFE: 18 Months, if stored in original, unopened
bag in a cool dry well ventilated place.**

**MAISHA RAFUNI: Miezi 18 ikihifadhiwa kwenya mfuko
wake halisi isiofunguliwa na
iliowekwa pahali pakavu na baridi.**

**In case of an emergency, call Toll free Nos:
(Piga simu isiyo na malipo wakati wa shida)
0800720021 and 0800730030 (24hrs)**

@ = Registered trademark of a Syngenta Group Company

**FORMULATED IN KENYA BY:-
Twiga Chemical Industries Ltd.
under license of Syngenta Agro AG,
Basle, Switzerland**

**DISTRIBUTOR (MSAMBASAJI) IN KENYA:-
TWIGA CHEMICAL INDUSTRIES LTD.
P.O. BOX 30172,
NAIROBI, KENYA
TEL: +254 20 3942000**

syngenta

NET CONTENT: 25 KG

UZITO KAMILI: KILO 25



JINSI YA KUTUMIA
Kwa matokeo bora nafaka zitiliwe dawa pale
zinapoendelea kutiwa ghalani. Uharibifu
ukiokolea na kushika sana si rahisi kuuzuia.
Ingawa ni muhimu kueneza dawa kwenye nafaka
zote, lakini kwa kiasi fulani ule uwezokawe wa
kufanya kazi kwa msambao wa mvuke husaidia
kukamilisha kazi yote. Mbegu za nafaka zilizo kwa
dawa hukingwa kwa muda usiopungua miezi sita.

VIPIMO VYA KUTUMIA
Mahindi yaliyo kwenye mabunzi
Ni vigumu kuyakagua mahindi yaliyo kwenye
mabunzi dhidi ya dudumuzi. Matokeo safi ya
kukabiliana na wadudu yanapatikana kwa
mahindi yaliyopukuswa.

Nafaka (mahindi, ngano na maharagwe)
Kwenye nafaka iliyopukuswa (mahindi, ngano,
maharagwe, nakadhalika), Changanya gramu 50
(jaza kipimo ulichopewa), kwa kila gunia moja ya
nafaka ukichanganya na kifaa cha koleo.
Vihwagilia ACTELLC SUPER nyinjine kwenye
sehemu ya nje ya gunia lenye nafaka ili
kupunguza uwezekano wa shambulizi mpya.

Kumwagika
Safisha pahali palipomwagika ukitumia
mchanga au vumbi wa mbao na uzike mbali na
vianzo vya maji.

Jinsi ilivyo
ACTELLC SUPER ni dawa ya wadudu
iliyotengenezwa kwa msingi wa madini ya chuma
hifadhiwa ghalani, hufanya kazi ikimwagiliwa
mwagiliwa moja kwa moja au kwa nguvu za
msambao wa mvuke wake. Ikitumiwa kwa
kuzingatia maelezo yake hakuna dhana
zitatokea. Nafaka ilipotumiwa dawa tayari
zaweza kuviwa na binadamu na mifugo bila
shida.

Kumbuka
Muundo huu wa ACTELLC SUPER utatoa kinga
ya wadudu aina ya Dudumuzi (Larger grain
borer). Kama mzuko mpya wa wadudu
unatarajiwa pata ushauri kutoka kwa wataalamu

TAHADHARI WAKATI WA MATUMIZI
Hatua za usalama
Dawa hii usiifanyie kazi iwapo daktari
amekushauri kutumia kikazi vitu vilivyo na
muundo wa silisi ya madini ya chuma. Usivute,
kunywa au kula chochote unapofanya kazi na
dawa hii.

Kwa usa lama unapoichanganya
Vumbi ya dawa yoweza kudhuru macho. Kiwa
imeingia machoni osha kwa maji mengi. Vaa nguo
za kujikinga na dawa

Kwa salama baada ya kutumia
Futa na kupunguza vumbi lote la dawa kutoka
ngozini mwako. Osha mwili kwa maji na sabuni.
Badilisha na kuzifua nguo za kazi. Choma au
kizike chombo kitupu cha dawa. Usikutumie tena.
Iki kutohatarisha samaki, usimwage dawa isiyo
takwa katika maji.

UWEKAJI
Dawa hifadhiwe katika viwekeo vyake asili
vivyofungwa vizuri, mahali pakavu na baridi, na
iliyofungwa mbali na watoto na vyakula vya
mifugo

UTUPAJI
Toboa pakiti au uchome kisha fukia mbali na njia
yapitapo maji. Usitumie pakiti kwa kazi ni nyinje
yoyote. Lowesha na kulikausha vumbi la
dawa iliyomwagika ivyo kwa kutumia unga wa mbao
uliyoweshwa kwa majimaji, kisha fagia uchafu na
uufukie mbali na njia za maji.

Uchafu kwa mazingira
ACTELLC SUPER yaweza kudhuru nyuki na
samaki Kamwe usitupe dawa yoyote au mkebe
wake katika mito au vidimbwi.

MAELEZO YA HUDUMA YA KWANZA
Ikitutwa pumzi: Unga wa ACTELLC SUPER
yaweza kusababisha hali ya kupiga chafya na
kukohoa iwapo utapewa mwanya wa kuingilia njia
za upumuaji. Mtu mwenye kuhusika haraka
lazima aingie sehemu atakopipatia hewa safi.
Ngozini: Mwili uliochafuliwa na dawa uoshwe
kwa maji na sabuni.
Machoni: Minkio machoni ioshewe kwa maji safi
kwa muda wa dakika 15 Muone daktari.
Kiwa imezwa: Endapo ACTELLC SUPER
imezwa usilazimishie mgonjwa kutapika.
Haraka mpeleke mgonjwa pamoja na kibandiko
hiki kwa daktari.

MAELEZO YA KUONDOA SUMU
Dalili za sumu: Hizi ni pamoja na kuchefuchefu,
kuumwa na kichwa.
kutoona vizuri, mbano wakifuani, ulgevu wa
viungo na maumivu ya tumbo. Hali ikiwa mbaya
zaidi huenda kukawa na tatizo la kupumua,
pengine itakuwa lazima kumpa hewa na kumsaidi
kupumua.
Piga simu isiyo na malipo wakati wa dharura:
0800720021 na 0800730030 (masaa 24)

Kinga: Atropine sulphate ACTELLC SUPER ni
dawa iliyotengenezwa kwa msingi wa pyrethroid
na ya fosforasi. Iwapo italiwa shaka kuwepo
kwa sumu, dawa ya Atropine itolewe mara moja
kwa kudungwa sindano ya mshipa na mtu
mwenye ujuzi ni kipimo cha mg 2-5 ikifuatiwa na
mg 2 kila nusu saa. Wakati huo huo dawa ya
Pralidoxime (p-25 au P-2m) itolewe kwa kipimo
cha gramu 1 kwa mililita 2-3 ya maji kwa sindano
inayodungwa kwenye mshipa. Zisitolewe dawa za
kutuliza. Rudia matumizi ya dawa kutegemea
maendeleo ya matibabu.

ILANI KWA MTUMIAJI
Dawa hii ni sharti kutumiwa kulingana na maagizo
yaliyo kwenye kibandiko. Ni hatia chini ya sheria
za madawa kutumia au kuhifadhi dawa hii katika
hali isiyo salama.

THIBITISHO
Dhamana ya muezaji ni maelezo yaliyo kwenye
kibandiko ambayo ni lazima yafuatwe kikamilifu.
Munuzu lazima akubali hatari zote zinazoweza
kutokana na matumizi ya dawa hii kwa binadamu
au mali na lazima aikubali dawa hii katika hali
iliyo.