

**READ LABEL AND ATTACHED PACKAGING LEAFLET BEFORE USING.**  
**SOMA MAELEZO YOTE KWENYE FURUSHI NA UKURASA ULIOBANDIKWA KWA FURUSHI.**

**Shelf Life:** 2 years from date of manufacture in original unopened containers.

**Maisha rafuni:** Miaka miwili kuanzia tarehe ya kutengenezwa ikiwa viwekeo vyake havijafunguliwa.

**In case of an emergency Call Toll free No.:**

**Kukiwa na jambo la dharura piga simu isiotozwa Nambari:**  
0800 720 021 / 0800 730 030 (24 hr).

® Registered Trademark  
of a Syngenta Group Company

**Manufacturer / Watengenezaji:**  
Syngenta Crop Protection AG  
Basle, Switzerland

**Distributor / Wauzaji:**  
**Syngenta East Africa Ltd.**  
P.O. Box 30393  
00100 Nairobi, Kenya  
Tel.: 254-0703018000/9000  
Fax: 254-020-3007885

MEMBER OF

AGROCHEMICAL  
ASSOCIATION  
OF KENYA



MEMBERSHIP NO. 225

L1017233 KENY/01U PPE 4062212

 **SCORE 250 EC**®

**syngenta**



Emulsifiable concentrate  
Fungicide for use against  
ringspot in carnations  
and botrytis in Roses;  
angular leaf spot on French  
beans and Ascochyta leaf spot on snow peas.

Huchanganyika na maji

Dawa nzuri ya kuangamiza na kuzuia ringspot  
kwenye Carnations, botrytis kwenye roses,  
angular leafspot kwenye maharagwe na  
Ascochyta leaf spot kwenye mbaazi.

Guarantee / Dhamana: Difenoconazole 250 g/l.

Registration No. / Nambari Ya Kusajiliwa  
PCPB(CR) 0947

**Net Content: 1 Litre**  
**Ujazo: Lita 1**



**KEEP LOCKED OUT OF REACH  
OF CHILDREN.  
FUNGIA MBALI NA WATOTO.**

**750ml**

**Commercial and Agricultural Class.**  
**Kundi la Biashara na Kilimo.**

**Date of manufacture/  
Tarehe ya kutengeneza:**

**Expiry date/Tumia kabla ya:**

**Batch no./Nambari ya kifurushi:**

approx. measure

**500ml**

**250ml**

Product names marked ® or ™, the ALLIANCE FRAME  
the SYNGENTA Logo and the PURPOSE ICON  
are Trademarks of a Syngenta Group Company

**Fungicide for use against ringspot in carnations and botrytis in Roses; angular leaf spot on French beans and Ascochyta leaf spot on snow peas.**

**Mode of Action**

Score 250 EC® is taken up by the plant and acts on the fungal pathogen during penetration and haustoria formation. It stops development of the fungi by interfering with the biosynthesis of sterols in cell membranes. Although the mode of action permits protective and curative use, it is recommended to apply the product early enough to avoid irreversible crop damage and build up of the disease.

**Spectrum of Activity**

The strength of Score 250 EC® is its excellent activity on ringspot in carnations and botrytis in Roses; angular leaf spot on French beans and Ascochyta leaf spot on snow peas.

**Mixing and treatment procedure**

Fill the spray tank with half the required volume of water and start agitation. Add the required amount of product and continue agitation while adding the rest of the water and during spraying.

**Timing of Application**

The first spray should be timed at the onset of disease development.

For ornamental diseases, a maximum of 3 time applications of Score 250 EC® are recommended as from the onset of disease. For most vegetable diseases, a maximum of 4 applications of Score 250 EC® are recommended as from the onset of disease. Use 15-20 ml per 20-litre-knapsack-sprayer (based on a water volume of 400 lt./ha).

From the point of view of anti-resistance strategy, the repeated application of straight difenoconazole should be avoided; alternations or mixtures with fungicide(s) with different modes of action are recommended.

**DIRECTIONS FOR USE**

Crop	Disease	Rate (lt./ha)	Interval (days)
Roses	Botrytis	0.75	10-14
Carnations	Ring spot	0.5	14
Vegetables			
French Beans	Leaf spot (Angular)	0.5	7-10
Snow Peas	Leaf spot (Ascochyta)	0.4-0.5	10-14

**Crop Tolerance**

When used as recommended, Score 250 EC® is well tolerated by most crops it is recommended to be used on. In case of doubt, check crop tolerance in a small area before spraying a large area.

**Spray volumes**

Score 250 EC® can be applied with any high or medium volume ground and aerial application equipment provided with sufficient agitation. Ensure that the spray equipment is properly calibrated. Spray volume can vary between 100 and 1000 lt./ha.

**Pre-harvest interval:**

French Beans, Snow Peas: 14 days.

**Re-entry interval:**

Roses and Carnations: 6hrs.

**Storage**

Store in original tightly closed container away from sun and damp. Keep under lock and key, away from children, animals and other unauthorized persons.

**Safety Aspects**

**Precautionary measures**

KEEP LOCKED UP OUT OF REACH OF CHILDREN and other unauthorised persons, and animals.

**Operator safety**

Strictly avoid all contact of concentrate or spray mixture with skin, eyes or clothing. Avoid breathing spray mist. Prolonged or repeated exposure to product may be harmful. DO NOT eat or smoke during work; wash hands and face thoroughly first. WEAR pro-

TECTIVE clothing, i.e., overalls, gloves, gumboots and face shield. When applying indoors, wear respiratory mask, hat and goggles. Immediately wash off accidental splashes of product and spray mixture. Change contaminated clothing immediately and launder before re-use. After work wash entire body thoroughly with soap and water and change to clean clothing.

**Product and Container disposal:** Dispose of surplus product on a landfill site approved for pesticides or bury in a safe place away from water sources. Dilute surplus application mixture, pour it broadcast on fallow land. Dispose of plastic containers in a landfill site or burn in a safe place. Do not re-use empty containers for any other purpose, especially for food, feed or storing drinking water.

**NB:** Always comply with local legal requirements.

**Spillage and Leakage**

Remove heavily contaminated soil layer and bury in a safe place away from water sources. Spillages: soak up with sand, sawdust, soil or an oil-binding agent. Collect up the contaminated material and dispose of in a safe place or put in a waste bin or a plastic bag, mark it and dispose of on a landfill site approved for pesticides. Treat spill area with plenty of water.

### Environment and Wildlife

Non-toxic to bees. Harmful to fish. Avoid contamination of water used for irrigation or domestic purposes. No risk is expected for birds and earthworms, due to the low toxicity of the compound to these organisms. Keep all livestock out of freshly treated area.

### FIRST AID INSTRUCTIONS

**SKIN CONTACT:** Remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water.

**EYE CONTACT:** Rinse eyes with clean water for several minutes. If poisoning is suspected, stop work immediately and call a physician. Show label to the physician. If inhaled move to clean fresh air. If swallowed, repeatedly administer medicinal charcoal in plenty of water. Seek medical advice immediately if a large volume of concentrate was ingested. Keep the patient in a well-ventilated room and strip off contaminated clothes.

**Note:** Never give anything orally to an unconscious person and never induce vomiting. If any indisposition occurs stop work immediately and call a physician.

### TOXICOLOGICAL INFORMATION

**Poisoning symptoms:** No case of human poisoning is on record.

**Antidote:** No specific antidote is known.

**Note to the doctor:** Apply symptomatic therapy.

**In case of an emergency Call Toll free No.:**

0800 720 021 / 0800 730 030 (24 hr).

### NOTICE TO USER

This pest control product is to be used only in accordance with the instructions on the label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

### WARRANTY

Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

### Dawa nzuri ya kuangamiza na kuzuia ringspot kwenye Carnations, botrytis kwenye roses, angular leafspot kwenye maharagwe na Aschochyta leaf spot kwenye mbaazi.

Score 250 EC® huingia ndani ya mimea kwa kupitia kwa matawi. Score 250 EC® inapendekezwa kunyunyizwa kabla ya mvua kunyesha na kabla ya ugonjwa kuwa mwingi.

**Jinsi ya kutumia:** Score 250 EC® ikitumiwa kwa njia inavyopendekezwa hapa huzuia na huangamiza kungu kwenye roses.

### MAPENDEKEZO YA KUTUMIA

Mmea	Ugonjwa	Dawa lt./ha	Siku za kurudia kunyunyiza
Roses	Botrytis	0.75	10-14
Carnations	Ring spot	0.5	14
Maharagwe	Leaf spot (Angular)	0.5	7-10
Mbaazi	Leaf spot (Aschochyta)	0.4-0.5	10-14

### Kiasi cha maji kinachohitajika:

Kwa trakta au bomba la kubeba mgongoni (knapsack), tumia lita 200 hadi 1000 kwa hekta moja. Kila mara changanya tu kiasi cha dawa kitakachoweza kutumiwa na kumalizika siku hiyo.

### Siku za kupita kabla ya kuvuna baada ya mnyunyizo

Maharagwe, Mbaazi: siku 14.

**Subiri:** Ingia sehemu iliyonyunyiziwa baada ya masaa 6.

**Kuhifadhi:** Hifadhi mahali pakavu mbali na vyakula, watoto na mifugo. Chumba chenye dawa kiwe kikavu na chenye hewa ya kutosha na daima kiwe kimefungwa.

### Tahadhari

HIFADHI MBALI NA WATOTO, mifugo na watu wasiohusika nayo.

### Usalama kwa wanyunyizaji

Usimeze Score 250 EC®. Usivute hewa iliyo na dawa. Chunga dawa isiingie kwenye macho, mwili au nguo ulizovaa. Usile, usinywe na usivute sigara wakati wa kunyunyiza dawa; na kabla ya kufanya hivyo oga mikono na uso. Vaa nguo za kazi yaani ovaroli, buti, kofia na glavu hasa wakati wa kukoroga na kunyunyiza dawa. Ukimwagikiwa na dawa mwilini au nguoni, osha mara moja. Baada ya kazi oga mwili wote na maji na sabuni. Safisha nguo za kazi kabla ya kuzitumia tena.

### Mazingira na wanyama wa pori

Score 250 EC® hudhuru samaki lakini haidhuru nyuki. Usimwage dawa ovyo ovyo mtoni au kwenye kisima. Wanyama wote wasikubaliwe kuingia mahali Score 250 EC® imenyunyizwa na usiokote au kuvuna chakula yoyote ya mifugo mahali imenyunyizwa dawa.

### **Dawa imalizikapo**

Zika kiasi kidogo kidogo cha dawa iliyobaki ndani ya mashimo mbali mbali katika sehemu zisizo na mimea au maji. Ongeza maji mengi kwa ile dawa iliyobaki ndani ya bomba na halafu umwage mbali na mimea. Safisha vizuri bomba baada ya kazi. Usitumie mikebe au mapipa ya dawa kwa kazi yoyote, na hasa kwa kuwekea chakula, malisho ya mifugo au maji.

### **Kuvuja na kumwagika kwa dawa**

Udongo uliomwagikiwa na dawa uchofwe vizuri na utupilie mahali salama mbali na vyanzo au njia za maji. Kusanya dawa iliyomwagika kisha ifukie au weka kwenye mfuko wa plastiki na ukazike sehemu iliyo mbali na vyanzo vya maji. Sehemu iliyomwagikiwa dawa isafishwe kwa maji mengi.

**Zingatia:** Dawa iliyomwagika isitumike tena.

**Kinga:** Hakuna kinga maalum. Matibabu ni vile atakavyoona daktari.

### **Kukiwa na jambo la dharuru piga simu isiotozwa Nambari**

0800 720 021 /0800 730 030 (masaa 24)

### **HUDUMA YA KWANZA**

Ukijisilihi sumu imekuingia, acha kazi mara moja na fuata huduma ya kwanza kama ifuatavyo. Endapo dawa imekumwagikia, vuanzo zilizochofuka, nawa haraka kwa maji mengi na sabuni se-

hemu zote za mwili zilizopata dawa. Kama dawa imeingia kwenye macho yasafishe kwa maji mengi safi kwa dakika 15. Kama umevuta utusitusi wa dawa, nenda kwenye sehemu yenye hewa nyingi. Endapo dawa imemezwa, mgonjwa apewe mkaa wa hospitali uliochanganywa na maji mengi. Daktari aitwe mara moja kama kiasi kikubwa cha dawa kimemezwa.

**Zingatia:** Mgonjwa aliyepoteza fahamu asilishwe au kunyeshwa kitu chochote kwa njia ya mdomo, na wala asitapishwe. PELEKA MJERUHI KWA DAKTARI.

### **ILANI KWA MTUMIAJI**

Dawa hii ni sharti kutumiwa inavyopendekezwa hapa. Unaweza kushtakiwa usipotumia na kuweka kwa njia ya sawa.

### **THIBITISHO**

Dhamana ya muuzaji ni maelezo yaliyo kwenye kibandiko ambayo ni lazima yafuatwe kikamilifu. Mnunuzi lazima akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii kwa binadamu au mali na lazima aikubali dawa hii katika hali hiyo.

NO TEXT HERE

GLUE PAGE

PAGE NUMBER SEQUENCING IS DETERMINED BY THE FINAL LEADING EDGE.  
RIGHT HAND STARTS FROM BACK OF BOOKLET. LEFT HAND STARTS FROM FRONT OF BOOKLET.  
FIRST (FRONT) PAGE IS ALWAYS FIXED ON TOP OF BOOKLET.

BASE

NON-PRINTABLE