

Withdrawal times

- For meat : 21 days, For milk : 7 days.

STORAGE

- Store in a cool dry place away from direct sunlight
- Store Below 30°C.
- Protect from Light .

Warning

Keep out of reach of children.
Avoid the introduction of contamination during use.

PACKING

Available in a 50ml and 100ml bottle.




REGISTERED AND DISTRIBUTED BY:
Twiga Chemical Industries Ltd.
P.O. Box 30172 -00100, NAIROBI, KENYA
Tel: +254 20 3942000, Fax: +254 20 3942405

TWIGACYCLINE®

OXYTETRACYCLINE 20% LA INJECTION

for Cattle, Sheep, Goats & Pigs



Kwa tiba ya mifugo

For
Veterinary
use only



REGISTERED AND DISTRIBUTED BY:
Twiga Chemical Industries Ltd.
P.O. Box 30172 -00100, NAIROBI, KENYA
Tel: +254 20 3942000, Fax: +254 20 3942405

TWIGACYCLINE

POM

**Oxytetracycline 20% LA Injection
in Cattle, Sheep, Goats & Pigs
FOR VETERINARY USE ONLY**

Description

Oxytetracycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria like *Bordetella*, *Campylobacter*, *Chlamydia*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. The action of oxytetracycline is based on inhibition of bacterial protein synthesis. Oxytetracycline is mainly excreted in urine, for a small part in bile and in lactating animals in milk. One injection acts for two days.

Composition

Contains per ml.:
Oxytetracycline base : 200 mg.
Solvents ad. : 1 ml.

Indications

Treatment of diseases caused by Oxytetracycline susceptible organism in cattle, camels, sheep and goats.

Diseases including pneumonia and shipping fever complex associated with *Pasteurella* spp. and *Haemophilus* spp. infectious bovine keratoconjunctivitis (pinkeye) caused by *Moraxella Bovis*, foot rot and diphtheria caused by *Fusobacterium Necrophorum*, bacterial enteritis (scours) caused by *Escherichia coli*, wooden tongue caused by *Actinobacillus lignieresii*, leptospirosis caused by *Leptospira pomona*. Wound infections and acute metritis caused by strains of staphylococci and streptococci organisms sensitive to Oxytetracycline.



Contra indications

- Hypersensitivity to tetracyclines.
- Administration to animals with a serious impaired renal and/or liver function.
- Do not use in cats, dogs and horses.
- Concurrent administration with penicillines, cephalosporines, quinolones and cycloserine.

Side effects

- After intramuscular administration local reactions can occur, which disappear in a few days.
- Discoloration of teeth in young animals.
- Hypersensitivity reactions.

DOSAGE AND ADMINISTRATION

Administer by deep intramuscular injection. The dose is 20mg/kg bodyweight (1ml/10kg bodyweight). The product is recommended for a single administration. Maximum recommended dose at one site:
This dosage can be repeated after 48 hours when necessary.

Do not administer more than 20 ml. in cattle, more than 10 ml. in swine and more than 5 ml. in calves, goats and sheep per injection site.