

TWIGACYCLINE 20% LA INJECTION

For Veterinary Use Only

WITHDRAWAL PERIOD

21 days for meat and edible tissues

7 days for milk

COMPOSITION

Each mL contains

Oxytetracycline 200mg.

STORAGE

Store in a cool dry place away from direct sunlight

PRESENTATION:

Available in a 100ml bottle.

MFG:

EXP:

Batch NO.

Manufactured by:
HEBEI YUANZHENG PHARMACEUTICAL
LIABILITY CO. LTD.
SHIJIAZHUANG, HEBEI, CHINA

A product of:
Twiga Chemical Industries Ltd.
P.O. Box 30172 -00100, NAIROBI, KENYA

TWIGACYCLINE

Oxytetracycline 20% LA Injection in Cattle, Sheep, Goats & Pigs

For Animal Treatment Only

Keep Out of Reach of Children

Store Below 30°C.

Protect from Light .

Avoid the introduction of
contamination during use.



100ML

INDICATIONS

Treatment of diseases caused by Oxytetracycline susceptible organism in cattle, camels, sheep and goats
Diseases including pneumonia and shipping fever complex associated with *Pasteurella spp.* and *Haemophilus spp.*, infectious bovine keratoconjunctivitis (pinkeye) caused by *Moraxella Bovis*, foot rot and diphtheria caused by *Fusobacterium Necrophorum*, bacterial enteritis (scours) caused by *Escherichia coli*, wooden tongue caused by *Actinobacillus lignieresii*, leptospirosis caused by *Leptospira pomona*. Wound infections and acute metritis caused by strains of staphylococci and streptococci organisms sensitive to Oxytetracycline.

DOSAGE AND ADMINISTRATION

Administer by deep intramuscular injection. The dose is 20mg/kg bodyweight (1ml/10kg bodyweight). The product is recommended for a single administration. Maximum recommended dose at one site:

Cattle	20ml
Sheep & Goats	5ml
Pigs	10 ml