

DOSAGE AND ADMINISTRATION

TWIGAMEC is to be given only subcutaneously. Inject under the loose skin in front of or behind the shoulder. Replace with a fresh sterile needle after every 10-20 animals.

The recommended dosage level is 200 mcg ivermectin per kg bodyweight. (1ml per 50kgs bodyweight). Use the following dosage schedule:

Bodywt (kg)	Dose Vol (ml)
Upto 50	1.0
51-100	2.0
101-150	3.0
151-200	4.0
201-250	5.0
251-300	6.0
301-350	7.0
351-400	8.0
401-450	9.0
451-500	10.0

Manufactured by: HEBEI YUANZHENG PHARMACEUTICAL LIABILITY CO. LTD. SHIJIANZHUANG, HEBEI, CHINA

A product of: TWIGA CHEMICAL INDUSTRIES LTD. P.O. BOX 30172 - 00100, NAIROBI, KENYA

TWIGAMEC

Ivermectin 1% Injection

For Treatment of Ecto & Endo parasites in Cattle, Sheep & Swine

POM-VPS

For Animal Treatment Only

Keep Out of Reach of Children

Store Below 30⁰C.

Protect from Light .

Avoid the introduction of contamination during use.



TWIGA
CHEMICALS

100ml



TWIGAMEC 1% INJECTION

For Veterinary Use Only

COMPOSITION

Each ml contains Ivermectin 10mg

WITHDRAWAL PERIOD

28 days for cattle, camel

21 days for pig, sheep, goat

21 days for milk

STORAGE

Store in a cool dry place, away from direct sunlight

PRESENTATION: TWIGAMEC

Injection is available in a 100ml vial.

MFG:

EXP:

Batch NO.