

TWIGAZOLE[®] 2.5%

ALBENDAZOLE 2.5% v/w DRENCH



Protects against round, tape worms & Liver flukes.



FOR VETERINARY USE ONLY

KEEP OUT OF REACH OF CHILDREN

5 Litre

TWIGAZOLE[®] 2.5%

ALBENDAZOLE 2.5% v/w DRENCH

DOSAGE

Dosage in Cattle		
BODY WEIGHT	Worm Dose (7.5 mg/kg)	Fluke & Worm (10 mg/kg)
50 Kg	15 ml	20 ml
100 Kg	30 ml	40 ml
150 Kg	45 ml	60 ml
200 Kg	60 ml	80 ml
250 Kg	75 ml	100 ml
300 Kg	90 ml	120 ml
350 Kg	105 ml	140 ml
400 Kg	120 ml	160 ml

Dosage in Sheep		
BODY WEIGHT	Worm Dose (5 mg/kg)	Fluke & Worm (7.5 mg/kg)
Up to 10 Kg	15 ml	20 ml
11 - 20 Kg	30 ml	40 ml
21 - 30 Kg	45 ml	60 ml
31 - 40 Kg	60 ml	80 ml
41 - 50 Kg	75 ml	100 ml
51 - 60 Kg	90 ml	120 ml
61 - 70 Kg	105 ml	140 ml
71 - 80 Kg	120 ml	160 ml

Albendazole 2.5% Each ml contains 25mg Albendazole

RECOMMENDATION

In sheep it is active against benzimidazole - susceptible strains of the following species:

Roundworms: Ostertagia, Haemonchus, Trichostrongylus, Nematodirus (including N. battus), Chabertia and Oesophagostomum. It is usually effective against inhibited larvae of Ostertagia.

Lungworms: Dictyocaulus filaria.

Tapeworms: Moniezia spp.

Adult Liver Fluke: Fasciola hepatica.

In cattle it is active against the following species:

Roundworms: Ostertagia, Haemonchus, Trichostrongylus, Nematodirus, Oesophagostomum, Bunostomum, Cooperia and Strongyloides spp. It is usually effective against inhibited larvae of Cooperia and Ostertagia.

Lungworms: Dictyocaulus viviparus.

Tapeworms: Moniezia spp.

Adult Liver Fluke: Fasciola hepatica.

- Withdrawal: Meat- 14 days, Milk 3 days.
- Shake well before use. Store below 25°C.
- Do not administer during first 3 weeks of pregnancy.

A product of: Twiga Chemical Industries Ltd.
P O BOX 30172-00100, NAIROBI-KENYA.
TEL: +254 20 3942000/ 3942300, FAX: 3942405

Batch No.:

Mfg Date:

Exp. Date: