




# TWIGAMEC

Ivermectin 1% Injection

**For Treatment of Ecto & Endo parasites in Cattle, Sheep & Swine**



**DISTRIBUTOR (MSAMBAZAJI) IN KENYA:-**  
Twiga Chemical Industries Ltd.  
P.O. Box 30172, 00100, GPO NAIROBI, KENYA  
Tel: 020-3942300, Fax: 3942405

***We Care for Your Livestock***

# TWIGAMEC

Ivermectin 1% Injection

## For Treatment of Ecto & Endo parasites in Cattle, Sheep & Swine

### PRESENTATION:

Twigamec is a clear, colourless, slightly viscous, Sterile solution containing 1% Ivermectin.

### Specific Indications:

Twigamec is indicated for the control of gastro-intestinal nematodes, lungworms, lice and mange mites.

**SPECIES:** Cattle, Sheep, Goats and Pigs

### DOSAGE & ADMINISTRATION:

#### Cattle:

**Dosage:** The recommended dose is 1 ml per 50 kg bodyweight.

**Administration:** Inject subcutaneously in front or behind the shoulder using aseptic technique.

#### Pigs:

**Dosage:** The recommended dose is 1 ml per 33 kg bodyweight.

**Administration:** Inject subcutaneously into the neck using aseptic technique.

### RECOMMENDED WITHHOLDING PERIOD:

#### Cattle:

- Cattle must not be slaughtered for human consumption within 21 days after treatment.
- Do not treat cattle producing milk for human consumption or in dairy cows within 60 days before calving.

#### Pigs:

- Pigs must not be slaughtered for human consumption within 21 days after treatment.

Bodywt (kg)	Dose Vol (ml)
Upto 50	1.0
51-100	2.0
101-150	3.0
151-200	4.0
201-250	5.0
251-300	6.0
301-350	7.0
351-400	8.0
401-450	9.0
451-500	10.0

### PHARMACEUTICAL PRECAUTIONS:

- Once vial is broached, the product should be used within 28 days.
- The product is extremely dangerous to fish and aquatic life.
- Do not contaminate surface waters or ditches with product container or unused product.

**PACKS:** 100ml, 50 ml and 10ml vials.

